



Toward elimination of mother-to-child transmission of HIV: The impact of a Rapid Results Initiative in Nyanza Province, Kenya




Ndege V.^{1,2}, Dillabaugh L.^{2,3}, Lewis-Kulzer J.^{2,3}, Owuor K.^{1,2}, Oyanga A.^{1,2}, Shade SB.^{2,3}, Akama E.^{1,2}, Agengo G.^{1,2}, Apuot S.^{1,2}, Otieno JP.^{1,2}, Onono M.^{1,2}, Odhiambo N.^{1,2}, Oyaro P.^{1,2}, Ngugi E.⁴, Bukusi E.A.^{1,2}, Cohen C.R.^{2,3}



1. Kenya Medical Research Institute (KEMRI), Nairobi, Kenya
2. Family AIDS Care and Education Services, Kisumu, Kenya
3. University of California San Francisco, San Francisco, CA, USA
4. Division of Global HIV/AIDS, U.S. Centers for Disease Control and Prevention, Kisumu, Kenya

Background

Despite extensive scale-up of Prevention of Mother-to-Child Transmission (PMTCT) services in Kenya, many HIV-infected women and exposed infants are not receiving comprehensive preventive services. Novel approaches are needed to improve PMTCT provision.



Methods

Rapid Results Initiative: A Rapid Results Initiative (RRI) was designed and implemented by Family AIDS Care and Education Services (FACES) and the Kenyan Ministries of Health to address key challenges including CD4 testing, highly active antiretroviral therapy (HAART) initiation for pregnant women and infants, male involvement, and early infant diagnosis (EID).




Figure 1. Rapid Results Initiative (RRI) Approach

Setting: This RRI intervention and evaluation was carried out at 119 FACES-supported health clinics in 5 districts in Nyanza Province, including: 6 district hospitals, 5 sub-district hospitals, 26 health centers, and 82 dispensaries.

Measurement: Site-level data were captured using routine monthly reports during the following 12-week timeframes:

Timeframe	Start	End
Baseline	October 2010	January 2011*
During the RRI intervention	April 2011	June 2011
Post RRI intervention	September 2011	July 2011

*Note: Dec. 2011 omitted due to the short work month

Analysis: Data obtained during the baseline, RRI and post-RRI periods were compared to assess whether there were significant changes during the three periods using pre-post cohort analysis using STATA 10. The risks, risk difference and risk ratios (95% Confidence Intervals) were reported for each indicator with the RRI baseline period as the reference point.



Photo by Jonathon Vlahos

Results

CD4 testing: CD4 testing uptake among HIV-infected pregnant women increased by 13% (RR=1.1, 95% CI=1.1-1.2) during RRI and remained elevated post-RRI (RR=1.07, 95%CI 1.02-1.1) compared to baseline.

Maternal HAART initiation: The relative proportion of HAART initiation improved from 13.7% to 19.7% among pregnant HIV-infected women (RR=1.4, 95% CI=1.2-1.7) during the RRI and to 21.7% (RR=1.6, 95% CI=1.4-1.8) post-RRI.

EID uptake: Uptake of EID among exposed infants increased by 30% during RRI (RR=1.3, 95% CI=1.2-1.4) and by 90% post-RRI (RR=1.9, 95% CI=1.8-2.0).

Infant HAART initiation: Infants initiated on HAART increased from 54.8% to 60.1% (RR=1.1, 95% CI=0.9-1.4) during RRI and to 69.0% post-RRI (RR=1.3, 95% CI=1.0-1.6).

Male partner engagement: Male partner testing increased from 7.7% at baseline to 16.4% during the RRI (RR=2.1, 95% CI=2.0-2.3). In the post-RRI period male partner testing decreased to 11.5% but remained 1.5 times above baseline figures (RR 1.5, 95%CI 1.4-1.7).

Table 1: Summary of Key Outcomes

HIV	Baseline Period Oct 2010 - Jan 2011*	RRI Period Apr - Jun 2011	Post - RRI Period Jul - Sep 2011
	N/%	N/% (95%CI)*	N/% (95%CI)*
Maternal outcomes			
HIV testing	8591	9123	8068
positive	1662	1890	1526
CD4 testing	59.0%	66.6% 1.1 (1.1-1.2)	63.3% 1.1 (1.0 - 1.1)
HAART initiation	13.7%	19.7% 1.4 (1.2 - 1.7)	21.7% 1.58 (1.4 - 1.8)
HIV - exposed infant outcomes			
PCR testing	46.2%	60.8% 1.3 (1.2 - 1.4)	87.0% 1.9 (1.8 - 2.0)
PCR positive	12.1%	13.8% 1.1 (0.9 - 1.4)	11.5% 0.9 (0.7 - 1.2)
HAART initiation	54.8%	60.1% 1.1 (0.9 - 1.4)	69.0% 1.3 (1.0 - 1.6)
Male partner engagement outcome			
HIV testing	7.7%	16.3% 2.1 (2.0 - 2.3)	11.6% 1.5 (1.4 - 1.7)

*December 2010 data excluded

*Results compared to baseline survey period

Bold - Statistically significant

Conclusions

Significant and sustained improvement in PMTCT services can be achieved using an RRI intervention. Similar strategies should be employed country-wide to work toward eliminating vertical transmission.

Literature cited

1. L. Mofenson, "Protecting the Next Generation – Eliminating Perinatal HIV-1 Infection", *N Engl J Med*, vol 362, no. 24, pp. 2316-2318, 2010.
2. P.M. Barker, W. Mphatswe, N. Rollins N. "Antiretroviral drugs in the cupboard are not enough: the impact of health systems' performance on mother-to-child transmission on HIV." *J Acquir Immune Defic Syndr*, vol. 56, no. 2, pp. 45-48, 2011.
3. National AIDS and STI Control Program (NASCOP), Kenya AIDS Indicator Survey KAIS 2007, Final Report, Ministry of Health, Kenya, 2009. http://nascop.or.ke/library/3d/Official_KAIS_Report_20091.pdf
4. Kenya National Bureau of Statistics (KNBS) and ICF Macro, Kenya Demographic and Health Survey 2008-2009. Calverton, Maryland: KNBS and ICF Macro, 2010.

Acknowledgments

We wish to acknowledge Family AIDS Care and Education Services (FACES), the University of California San Francisco (UCSF), the Kenya Medical Research Institute (KEMRI), the Kenyan Ministries of Health (MOH), U.S. President's Emergency Plan for AIDS Relief (PEPFAR) and the U.S. Centers for Disease Control and Prevention (CDC), and the women, men and children in the communities we serve.

Sources of Support

FACES is a UCSF and KEMRI collaborative program that is supported by PEPFAR/CDC(Cooperative Agreement 1U2GPS001913).

Disclaimer

The content and views in this abstract are solely the responsibility of the authors and do not necessarily represent the official views of the affiliated organizations, United States Government, or Government of Kenya.



Presented at AIDS 2012 – Washington, D.C., USA

