

Extended Nevirapine Prophylaxis for Infants and Its Impact on Vertical Transmission Rates at 18 months, a Retrospective Cohort Study

Akama, E.¹, Oyaro, P.¹, Nimz, A.², Mburu, M.²,
Diouf, K.³, Hongo B.⁶, Lewis-Kulzer J.^{1,5},
Cohen, C.R.^{1,5}, Bukusi, E.A.,¹, Abuogi, L.^{1, 4}

Kenya Medical Research Institute, Research Care and Training Program, Family AIDS Care and Education Services (FACES), Kisumu, Kenya¹

University of Colorado School of Medicine², Department of Pediatrics⁴, Aurora, Colorado

Department of Obstetrics, Gynecology and Reproductive Biology, Division of Global OB GYN, Brigham and Women's Hospital, Boston, MA, USA³

University of California San Francisco, Department of Obstetrics, Gynecology & Reproductive Sciences⁵

Ministry of Health⁶

BACKGROUND

- Interventions to prevent mother to child transmission (MTCT) of HIV in many settings are complicated by necessarily high rates of prolonged breastfeeding
- In sub-Saharan Africa up to 15% of HIV infections in infants may be attributed to breastfeeding

OBJECTIVES

- To determine the rate for MTCT in infants on extended nevirapine (eNVP) prophylaxis during breastfeeding at eighteen months of age

METHODS

Setting

- FACES is a PEPFAR-funded comprehensive HIV prevention, care and treatment program based in Migori, Homa bay, Kisumu Counties
- The study was conducted in FACES supported Kenyan Ministry of Health (MOH) clinics :
 - Lumumba Health Center, Kisumu County
 - Migori District Hospital, Migori County
 - Oyani Health Center, Migori County

METHODS

Time Interval:

- eNVP Cohort – June 2011 to November 2013
- Historical Cohort – January 2008 to January 2011

This cohort study compared:

- MTCT rates between a group of prospectively followed infants receiving eNVP prophylaxis
- A historical cohort who received single dose NVP at birth with 6 weeks of zidovudine (AZT)



METHODS

Table 2. Daily nevirapine (dNVP) prophylaxis for HIV-exposed infants

Age	Nevirapine dose
0 – 6 weeks	Birth weight <2500 g – 10 mg (1 ml) once daily Birth weight >2500 g – 15 mg (1.5 ml) once daily
6 weeks – 14 weeks	20 mg (2 ml) once daily
14 weeks to 6 months	25 mg (2.5 ml) once daily
6 months – 9 months	30 mg (3 ml) once daily
9 months – 12 months	40 mg (4 ml) once daily
>12 months	50 mg (5 ml) once daily

METHODS

Analysis

- Data was transferred to STATA 12 for all the analysis
- Bivariate analysis was done to determine significant differences between cohorts and infection rate using Chi Square and Fishers Exact Tests as Appropriate
- A p-value < 0.05 was considered significant

RESULTS

Variable	Historical n(%) n=362	eNVP n(%) n=283	p-value
Results at week 6			
Negative	280(90.6)	274(97.9)	<0.001*
Positive	29(9.4)	6(2.1)	
Results at Month9			
Negative	205 (96.2)	213 (98.2)	0.256
Positive	8(3.8)	4(1.84)	
Results at 18 Months			
Negative	143(96.6)	172(100)	0.020*
Positive	5(3.4)	0(0)	
Cumulative Infection			
Negative	143(77.7)	172(94.5)	<0.001*
Positive	41(22.3)	10(5.5)	

CONCLUSION

- Implemented within a MOH PMTCT program:
 - Infant eNVP prophylaxis during the breastfeeding period, combined with appropriate ART for the mother during pregnancy, is associated with a low risk of HIV transmission
- Extended nevirapine (NVP) can be safely and effectively implemented in PMTCT programs

ACKNOWLEDGEMENTS

- Kenyan Ministries of Health (MOH)
- Family AIDS Care and Education Services (FACES)
- Kenya Medical Research Institute (KEMRI)
- University of California San Francisco (UCSF)
- U.S. President's Emergency Plan for AIDS Relief (PEPFAR)
- U.S. Centers for Disease Control and Prevention (CDC)
- The women, men and children in the communities served

The findings and conclusions in this presentation are those of the author(s) and do not necessarily represent the official position of U.S. Centers for Disease Control and Prevention/the and the Government of Kenya

This research has been supported by the President's Emergency Plan for AIDS Relief (PEPFAR) through the U.S Centers for Disease Control under the terms of Cooperative Agreement # PS001913

University of California
San Francisco

