


# Migrant fisherfolk and factors associated with HIV clinic attendance in Suba District, Kenya

Lindah Otieno<sup>1,2</sup>, Kevin Owuor<sup>1,2</sup>, Jeremy Penner<sup>2,3</sup>, Jayne Lewis Kulzer<sup>2,4</sup>, Starley Shade<sup>2,4</sup>, Dickens Otieno<sup>1,2</sup>, John Owiti<sup>1,2</sup>, Craig R. Cohen<sup>2,4</sup>, Elizabeth A. Bukusi<sup>1,2</sup>

1. Kenya Medical Research Institute (KEMRI), Nairobi, 2. Family AIDS Care and Education Services, Kisumu, 3. University of British Columbia, Vancouver, 4. University of California San Francisco, San Francisco

## Background

- ◆ Suba – an isolated island district with a migrant fishing population
- ◆ Strategies have been implemented to increase HIV service access and retention for hard-to-reach fisherfolk:
  - ➔ Static mobile HIV clinics on the island and a toll-free hotline to organize medication pick-ups and clinic visits have been established



## Setting

- ◆ Family Aids Care and Education Services (FACES) is a HIV prevention, care, and treatment program
  - ➔ Collaborative University of California San Francisco (UCSF) and Kenya Medical Research Institute (KEMRI) program
  - ➔ Funded by U.S. President's Emergency Plan for AIDS Relief (PEPFAR) Centers for Disease Control and Prevention (CDC)
- ◆ FACES supports HIV services in all health facilities in the island and two mobile clinics.
- ◆ Fisherfolk definition:
  - ➔ Fishermen
  - ➔ Boat owners
  - ➔ Fishmongers
  - ➔ 'Abila' – women who cook for the fishermen



## Methods

- ◆ Retrospective cohort evaluation conducted to determine risk factors for HIV care defaulting among fisherfolk at three health facilities
  - ➔ One static clinic: Sena Health Centre
  - ➔ Two mobile clinics: Remba and Ringiti Health clinics
- ◆ Inclusion criteria
  - ➔ Patients who indicated fisherfolk as their occupation
  - ➔ Enrolled in HIV care between January 2009 and September 2010
- ◆ Nine months later, study participants files were reviewed to determine
  - ➔ Defaulting status (missed appointment by >3 days)
  - ➔ Potential risk factors: age, gender, CD4 count, WHO stage, anti-retroviral (ART) status, and disclosure
- ◆ Multivariate logistical regression analysis was performed to calculate odds ratios (OR) and 95% confidence intervals (CI)

## Results

Among 418 fisherfolk files examined:

- ◆ Age: Mean 30.4 years (SD; 11.3)
- ◆ Gender: 268 (64%) female
- ◆ ART status: 322 (77%) on ART
- ◆ Youth Status: 24 (6%) youth (ages 14-21 years)
- ◆ Default Events: Median 1 (IQR; 0,2)
- ◆ Defaulting Proportion: 251 (60%) defaulted at least once

Table 1: Multivariate analysis of predictors of defaulting

Variables	Multivariate Analysis		
	OR	95% CI	P Value
Non-Disclosure	6.4	2.9-14.1	<0.001
Youth (14 – 21yrs)	3.9	1.2-12.6	0.021
CD4 = 350	2.8	1.4-5.5	0.002

\*Not associated with defaulting: Gender, WHO stage, and ART status

## Limitations

- ◆ The risk of defaulting among the fisherfolk was not compared with the non- fisherfolk in the island population
- ◆ Marital status of the fisherfolk was not examined

## Conclusion

- ◆ Fisherfolk were more likely to default from HIV care if they had not disclosed their status, were young, or had higher CD4 counts
- ◆ Interventions to improve disclosure warrant evaluation and HIV programs require targeted interventions to support youth and retention efforts that extend beyond the sickest patients



## Acknowledgment

- ◆ U.S. President's Emergency Plan for AIDS Relief (PEPFAR)
- ◆ U.S. Centers for Disease Control and Prevention (CDC)
- ◆ Family AIDS Care and Education Services (FACES)
- ◆ Kenya Medical Research Institute (KEMRI)
- ◆ Ministry of Public Health & Sanitation
- ◆ Ministry of Medical Services
- ◆ Students from UCSF (Michelle, Diana & Teja)
- ◆ Abasuba community